

**MYOCARDIAL ISCHEMIA AND INFARCTION**

**C-TERMINAL PROVASOPRESSIN (COPEPTIN), MID-REGION PRO-ADRENOMEDULLIN (MR-PROADM) AND MID-REGION PRO-ATRIAL NATRIURETIC PEPTIDE (MR-PROANP) AS NOVEL PROGNOSTIC BIOMARKERS AFTER NON-ST ELEVATION ACUTE CORONARY SYNDROME: ANALYSIS FROM MERLIN TIMI-36**

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Monday, April 04, 2011, 9:30 a.m.-10:45 a.m.

Session Title: Unstable Ischemic Syndrome -- Clinical: Biomarkers

Abstract Category: 2. Unstable Ischemic Syndrome—Clinical

Session-Poster Board Number: 1071-330

Authors: Ryan G. O'Malley, Marc P. Bonaca, Marc S. Sabatine, Benjamin M. Scirica, Sabina A. Murphy, Petr Jarolim, Eugene Braunwald, David A. Morrow, TIMI Study Group, Brigham and Women's Hospital, Boston, MA

**Background:** Copeptin, MR-proADM, and MR-proANP are novel biomarkers of hemodynamic stress that have prognostic value in pts with heart failure (HF). We investigated the prognostic performance of these 3 biomarkers relative to established predictors in a large cohort with non-ST elevation acute coronary syndrome (NSTEMI-ACS).

**Methods:** We measured copeptin, MR-proADM, and MR-proANP at baseline in 4,282 pts with NSTEMI-ACS randomized to ranolazine or placebo in the MERLIN-TIMI 36 trial. Pts were followed for an average of 1 year. Cardiovascular death (CVD) and new or worsening HF were centrally adjudicated.

**Results:** High concentration (Q4 v. Q1-3) of each biomarker identified pts with an increased risk of CVD or HF at 12 months (Copeptin 13.2% v. 5.0%,  $p < 0.001$ ; MR-proADM 15.8% v. 4.1%,  $p < 0.001$ ; MR-proANP 17.7% v. 3.5%,  $p < 0.001$ ), as well as CVD and HF individually (Figure). After adjustment for important clinical covariates, each marker remained significantly associated with CVD and HF (Figure top row). Moreover, in a model that also included all 3 biomarkers plus BNP and troponin, a high concentration of each biomarker independently predicted an increased risk of CVD or HF (Copeptin HR 1.44, CI 1.12-1.83; MR-proADM HR 1.54, CI 1.16-2.04, and MR-proANP HR 1.67, CI 1.21-2.30), compared with BNP (HR 1.59, CI 1.15-2.20) with an increase in the c-statistic ( $p < 0.001$ ).

**Conclusion:** Copeptin, MR-proADM, and MR-proANP add to BNP, troponin, and clinical risk indicators for predicting CVD or HF in pts with NSTEMI-ACS.

**Event Rates at 12 months and Adjusted Hazard Ratios**

